

Neuropsychology

Altered Implicit Category Learning in Anorexia Nervosa

Megan E. Shott, J. Vincent Filoteo, Leah M. Jappe, Tamara Pryor, W. Todd Maddox, Michael D. H. Rollin, Jennifer O. Hagman, and Guido K. W. Frank

Online First Publication, December 26, 2011. doi: 10.1037/a0026771

CITATION

Shott, M. E., Filoteo, J. V., Jappe, L. M., Pryor, T., Maddox, W. T., Rollin, M. D. H., Hagman, J. O., & Frank, G. K. W. (2011, December 26). Altered Implicit Category Learning in Anorexia Nervosa. *Neuropsychology*. Advance online publication. doi: 10.1037/a0026771

Altered Implicit Category Learning in Anorexia Nervosa

Megan E. Shott
University of Colorado Denver

J. Vincent Filoteo
University of California San Diego

Leah M. Jappe
University of Minnesota

Tamara Pryor
Eating Disorder Center of Denver, Denver, Colorado

W. Todd Maddox
University of Texas

Michael D. H. Rollin, Jennifer O. Hagman, and
Guido K. W. Frank
University of Colorado Denver

Objective: Recent research has identified specific cognitive deficits in patients with anorexia nervosa (AN), including impairment in executive functioning and attention. Another such cognitive process, implicit category learning has been less studied in AN. This study examined whether implicit category learning is impaired in AN. **Method:** Twenty-one women diagnosed with AN and 19 control women (CW) were administered an implicit category learning task in which they were asked to categorize simple perceptual stimuli (Gabor patches) into one of two categories. Category membership was based on a linear integration (i.e., an implicit task) of two stimulus dimensions (orientation and spatial frequency of the stimulus). **Results:** AN individuals were less accurate on implicit category learning relative to age-matched CW. Model-based analyses indicated that, even when AN individuals used the appropriate (i.e., implicit) strategy they were still impaired relative to CW who also used the same strategy. In addition, task performance in AN patients was worse the higher they were in self-reported novelty seeking and the lower they were in sensitivity to punishment. **Conclusions:** These results indicate that AN patients have implicit category learning deficits, and given this type of learning is thought to be mediated by striatal dopamine pathways, AN patients may have deficits in these neural systems. The finding of significant correlations with novelty seeking and sensitivity to punishment suggests that feedback sensitivity is related to implicit learning in AN.

Keywords: implicit, learning, reward, anorexia nervosa, temperament

Anorexia nervosa (AN) is a severe psychiatric illness characterized by restrictive eating, fears of being fat despite life threatening low weight, and can be associated with binge eating and/or purging behavior (APA, 2000). Common temperament traits of eating disorder patients include high harm avoidance, cognitive rigidity, and alterations in feedback sensitivity (Anderluh, Tchanturia, Rabe-Hesketh, & Treasure, 2003; Bulik et al., 2003; Jappe et

al., 2011; Wagner et al., 2006). AN has the highest mortality among all psychiatric disorders (Sullivan, 1995), and as such, understanding this disease from multiple perspectives is an important undertaking.

In recent years, there has been growing interest in studying cognitive functioning in patients with AN. These studies are important for a better understanding of AN given the possibility that cognitive deficits may (a) contribute to the development and persistence of AN, (b) result from neurological changes associated with the disease, or (c) influence the choice of treatment approaches. Many of these studies have found deficits primarily in executive functioning (Fassino et al., 2002; Holliday, Tchanturia, Landau, Collier, & Treasure, 2005; Jones, Duncan, Brouwers, & Mirksy, 1991; Lauer, Gorzewski, Gerlinghoff, Backmund, & Zihl, 1999; Roberts, Tchanturia, Stahl, Southgate, & Treasure, 2007; Steinglass, Walsh, & Stern, 2006; Tchanturia, Whitney, & Treasure, 2006), visual-spatial abilities (Grunwald et al., 2001; Jones et al., 1991; Kingston, Szmukler, Andrewes, Tress, & Desmond, 1996; Palazidou, Robinson, & Lishman, 1990; Szmukler et al., 1992), and attention (Bosanac et al., 2007; Johansson, Ghaderi, & Andersson, 2005; Ohrmann et al., 2004; Seed, Dixon, McCluskey, & Young, 2000; Seed, McCue, Wesnes, Dahabra, & Young, 2002; Shafraan, Lee, Cooper, Palmer, & Fairburn, 2007). Although past

Megan E. Shott, Michael D. H. Rollin, and Jennifer O. Hagman, Department of Psychiatry, University of Colorado, Denver; J. Vincent Filoteo, Department of Psychiatry, University of California, San Diego; Leah M. Jappe, Department of Psychology, University of Minnesota; Tamara Pryor, Eating Disorder Center of Denver, Denver, Colorado; W. Todd Maddox, Department of Psychology, University of Texas; Guido K. W. Frank, Departments of Psychiatry and Neuroscience, University of Colorado, Denver.

Correspondence concerning this article should be addressed to Guido K. W. Frank, MD, Assistant Professor, Departments of Psychiatry and Neuroscience, Director, Developmental Brain Research Program, University of Colorado Denver, The Children's Hospital, Gary Pavilion A036/B-130, 13123 East 16th Avenue, Aurora, CO 80045, Tel: 720-777-1909. E-mail: guido.frank@ucdenver.edu

studies have also identified deficits in AN patients in general intellectual functioning (Chui et al., 2008; Mathias & Kent, 1998), a recent systematic review and meta-analysis found that people with AN tend to have higher IQ scores compared with healthy comparison groups (Lopez, Stahl, & Tchanturia, 2010). These results suggest that difficulties in cognitive functioning in AN may be more specific than an impairment in general intellectual functioning, and raises the possibility that AN might impact only certain neural systems. In a recent review, Lena et al. (Lena, Fiocco, & Leyenaar, 2004) stressed the need for greater exploration of these neurocognitive deficits in AN suggesting that impaired cognition may in fact contribute to the genesis of the disease.

Although there have been more recent studies examining cognition in AN, there are other potentially important processes that have received less attention, such as those associated with learning and memory (Chui et al., 2008; Jones et al., 1991; Mathias & Kent, 1998). This area of research could have important implications for a better understanding as to whether or how faulty learning and memory mechanisms contribute to the development or maintenance of AN. While a handful of studies have examined explicit memory performance in AN, few have studied implicit memory in these patients. Implicit memory denotes a class of learning and memory processes that operate outside conscious awareness and includes processes such as priming, classical conditioning, and procedural learning. The few studies that have studied implicit learning in AN have examined whether patients with an eating disorder are abnormally primed to process food- or body-related information (Hermans, Pieters, & Eelen, 1998; Johansson, Ghaederi, Hallgren, & Andersson, 2008). The results of those studies have been somewhat inconsistent. Whereas Hermans et al. (1998) found no implicit memory bias in their word stem completion task, Johansson et al. (2008) found that AN individuals exhibited attentional interference for food-related words. Other forms of implicit learning that have been examined include measures of procedural learning, which is a process primarily involved in the acquisition of skills and habits. Lawrence et al. (Lawrence et al., 2003), for example, found that AN patients were impaired relative to controls on a measure of visual discrimination learning (which is believed to be a measure of procedural learning). These same patients were not impaired on a pattern recognition task, which is a measure of explicit memory. These authors argued that the AN patients' deficits on the visual discrimination learning task were likely related to alterations in the subcortical dopamine reward-based learning system, given that alterations in this system in primates and humans diminishes the learning of this task. In contrast, the normal performance of the AN patients on the pattern recognition task was thought to be related to intact functioning in the medial temporal lobe memory system, which is thought to mediate explicit learning and memory. The suggestion that changes in the dopamine reward-based learning system resulted in AN patients' deficits on this task is consistent with a growing literature suggesting alterations in the dopamine system in these patients (Fladung et al.; Frieling et al., 2010; Kaye, Ebert, Gwirtsman, & Weiss, 1984).

Additional evidence that AN patients may have alterations in reward-based mechanisms comes from studies using self-reports, such as the Sensitivity to Punishment-Sensitivity to Reward Questionnaire (SPSRQ), which measures the saliency of rewarding or punishing real-world reinforcers (e.g., social or monetary reinforc-

ers). Using the SPSRQ, Jappe et al. (2011) found that AN patients were more sensitive to both punishment and reward than controls, indicating alterations in feedback sensitivity in AN patients. Despite these interesting findings, the relationship between the AN patients' altered feedback sensitivity and behavioral measures of reinforcement-based learning have yet to be examined. The potential link between learning and feedback sensitivity in AN is important in this context, since traits such as sensitivity to reward or punishment affect learning (Cavanagh, Frank, & Allen, 2011) and could be related to the ability to change behavior and thus the course of the illness. Taken together, the study of implicit learning, and how deficits in this area relate to self-report AN temperament or behaviors, may help identify specific brain alterations in AN, which could have important implications in our understanding of what AN individuals learn to find rewarding or reinforcing (Wächter, Lungu, Liu, Willingham, & Ashe, 2009).

The purpose of the present study was to examine implicit learning in women with AN. To this end, AN patients and control women (CW) were administered a category learning task that emphasized the procedural-based learning system that relies on dopamine reward-based learning (Ashby, Alfonso-Reese, Turken, & Waldron, 1998). The task is described in detail below, but briefly, participants are presented with simple perceptual stimuli, categorize the stimuli into one of two categories, and are given corrective feedback immediately following their response. Importantly, this task is believed to be mediated by the dopamine reward-based learning system since fast, corrective feedback stimulates dopamine activation and learning (Schultz, 2002), while delaying such feedback results in diminished learning (Maddox, Ashby, & Bohil, 2003; Maddox et al., 2005). Other evidence that dopamine signaling might play an important role in learning the procedural-based category learning task comes from studies showing that patients with known damage to the dopamine systems (i.e., Parkinson's disease) are impaired on this task (Filoteo, Maddox, Salmon, & Song, 2005) as well as functional neuroimaging studies with normal participants demonstrating activation of the posterior caudate (Nomura et al., 2007), a brain region believed to make use of dopamine signaling during learning (Miyachi, Hikosaka, Miyashita, Karadi, & Rand, 1997; Nakamura & Hikosaka, 2006).

Given the previously identified alterations in dopamine systems in AN and the proposed involvement of such systems in implicit category learning, we predicted that AN patients would be significantly impaired in implicit learning relative to controls as demonstrated by lower accuracy on the category learning task. To assess whether implicit category learning was associated with temperament and behavior variables, we also examined the relationship between task performance and self-report measures of feedback sensitivity such as reward and punishment. We predicted that poorer performance in category learning would be associated with altered levels of feedback sensitivity in the AN group.

Method

Participants

Twenty-one female patients with AN and 19 control women (CW), all Caucasian, participated in the study. Ten of the AN patients were of the binge/purge subtype, while the remaining 11 AN patients were restricting subtype. Table 1 shows the mean age

Table 1
Demographic and Disease-Related Data for Study Groups

	CW (<i>n</i> = 19)		AN (<i>n</i> = 21)	
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>
Study Age (years)	27.3	5.3	25.2	6.4
BMI	21.8	1.5	16.2	1.3
BMI Low	—	—	14.4	2.3
Duration of Illness (years)	—	—	7.95	7.4
Age of Onset (years)	—	—	16.8	2.0

and body mass index (BMI) of the AN patients and CW participants, and the lowest BMI, duration of illness, and age of disease onset for the AN patients. The mean age of the AN group and the CW group did not differ ($p = .28$). As expected, the mean BMI for the AN patients was significantly lower than for CW participants ($p < .001$). A χ^2 test revealed that there were no educational level differences between the AN and CW groups ($p = .27$). Patients were recruited from the Partial Hospitalization Program at the Eating Disorder Center of Denver (a facility that treats adult eating disorder patients, $n = 17$) or the inpatient eating disorders unit at Children’s Hospital Colorado (a facility that treats children and adolescents with eating disorders through the age of 21, $n = 4$). While the mean age and duration of illness differed in the two treatment settings, AN patients had similar current or lowest BMI, and age of disease onset regardless of the type of treatment facility. Each patient completed the study within a few weeks of starting treatment and the subject’s height and weight were taken the day of testing. Patients completed the Structural Clinical Interview for *DSM-IV* Disorders (SCID-IV) with a trained doctoral level clinician to ensure they met criteria for AN. Control participants also completed the SCID to screen for abnormal eating behaviors or other *DSM-IV* disorders. All CW participants had a BMI between 19 and 25 and had no personal or family history of mental or neurological disorders. This study was approved by the Colorado Multiple Institutional Review Board and participants gave written informed consent.

Clinical Measures

The Beck Depression Inventory (BDI-2; Beck, Ward, Mendelson, Mock, & Erbaugh, 1961) is a 21-item self-report that measures depression symptom severity.

The Temperament and Character Inventory (Cloninger, Svrakic, & Przybeck, 1993) is a 240-item inventory consisting of 7 independent dimensions. We included two of these (Novelty Seeking and Harm Avoidance), which are variables previously implicated in AN (Wagner et al., 2006).

The Eating Disorder Inventory 2 (EDI-2; Garner, 1991) contains 91 items organized onto 12 primary scales. Of the 12 scales, four were used: Drive for Thinness, Body Dissatisfaction, Bulimia, and Perfectionism.

The Sensitivity to Punishment and Sensitivity to Reward Questionnaire (SPSRQ; Torrubia, Avila, Molto, & Caseras, 2001) is a 48-item self assessment that is based on Reinforcement Sensitivity Theory (RST; Gray, 1970). The scale provides separate scores for sensitivity to punishment (SP) and sensitivity to reward (SR).

Stimuli and Stimulus Generation

The specific version of the category learning task used in this study required participants to learn to categorize simple perceptual stimuli (Gabor patches; see Figure 1) into one of two categories (Category A or Category B) when the rule that dictated category membership was nonverbal (implicit learning) and required a linear integration of two stimulus dimensions (orientation and spatial frequency of the gratings of the Gabor patch; see below for details). The study used the randomization technique introduced by Ashby and Gott (Ashby & Gott, 1988). Stimuli consisted of Gabor patches that varied in spatial frequency and orientation (see Figure 1). For each experiment, an equal number of Category A and Category B stimuli were generated by sampling randomly from two bivariate normal distributions. Each category distribution is specified by a mean and a variance on each dimension, and by a covariance between dimensions. For both category structures it was always the case that the covariance matrix for Category A was identical to the covariance matrix for Category B. The categories differed only in the location of their means. The exact parameter values for each experiment are listed in Table 2 (Note: these are in arbitrary units that were transformed into the stimulus units using the methods described below). The category structure for the experiment is displayed in Figures 2. Each filled square in these figures denotes the spatial frequency and spatial orientation of a Gabor pattern from Category A, while each unfilled circle denotes the spatial frequency and spatial orientation of a Gabor pattern from Category B. The solid line in these figures denotes the location of the optimal decision bound. The use of the optimal bound maximizes long-run accuracy.




Figure 1. Computer generated Gabor patch.

Table 2
*Category Distribution Parameter Values for the Implicit
 Category Learning Task*

Category	Information-Integration (Implicit) Task				cov _{f,o}
	μ_f	μ_o	σ_f^2	σ_o^2	
A	260	125	75	9000	0
B	339	125	75	9000	0

The stimuli were computer generated and displayed on a 21 monitor with 1360×1024 resolution. In each of the three experiments, the stimuli consisted of a single Gabor patch (see Figure 1). Each of the stimuli varied in orientation and spatial frequency. Each Gabor patch was generated using MATLAB routines from Brainard's (1997) Psychophysics Toolbox. Each random sample (x_f, x_o) was converted to a stimulus by deriving the frequency, $f = .0025 + (x_f/5000)$ cycles per pixel, and orientation, $o = 0.36x_o$ degrees. The scaling factors were chosen in an attempt to equate the salience of frequency and orientation based on our past experience with these stimuli. Each Gabor patch was 7 cm in diameter, which subtended a visual angle of about 8.8 degrees from a viewing distance of 45 cm.

Experimental Procedure

Three hundred and twenty trials were presented and were broken down into four blocks of 80 trials. At the start of the experiment, the participants were told that they were involved in a study that examined their ability to categorize simple stimuli, that a series of stimuli would be presented, and that they would be asked to categorize each as a member of either Category A or Category B. They were also told that at the beginning of the experiment they may feel as though they were guessing, but as the experiment progressed, their accuracy would likely increase. Participants indicated their categorization responses by pressing one key for Category A stimuli and another key for Category B stimuli. For each trial, the stimulus was presented until the participant's categorization response was made, then immediately following their response, they were given feedback for 1 second that consisted of the word "wrong" if their response was incorrect or "correct" if their response was correct. Once feedback was given, the next trial was initiated 1 second later. Total task duration was between 20 and 25 minutes, depending on individual subject response time.

Model-Based Analyses

A major advantage of the task used in this study is that mathematical models have been developed to determine the specific approach used by participants when learning the task (Maddox & Ashby, 2004; Maddox et al., 2003; Zeithamova & Maddox, 2006). These models enable a more detailed investigation of procedural learning in that previous studies have shown that patient groups can differ from controls in implicit learning not because of an impairment in such processes, but instead because they adopted a different strategy than controls when learning (Shohamy, Myers, Onlaor, & Gluck, 2004). Thus, through the use of these models, we are able to identify AN patients who adopted a procedural-based

approach to learning compared to control individuals to determine if in fact the patients are impaired in procedural-based learning. Although the main dependent measure in this study was the accuracy of participants' responses, we also applied a series of models to the data to better understand the category learning processes used by the AN patients and CW participants. These models have been developed to analyze data in this paradigm and have been used extensively to better understand the underlying processes in category learning in both normal and patient populations (Ashby & Maddox, 1992; Ashby, 1992; Filoteo & Maddox, 1999; Filoteo, Maddox, & Davis, 2001; Maddox, 2001; Maddox & Ashby, 1993; Maddox & Ashby, 1996; Maddox & Filoteo, 2001; Maddox, Filoteo, Delis, & Salmon, 1996; Maddox, Filoteo, & Huntington, 1998). Two classes of models were applied to the final block of data separately for each participant (the details of these models can be found in (Maddox & Ashby, 1993; Maddox, 1999). One class of models assumes that the participant used a *procedural-based* (PB) approach by performing an implicit integration of the spatial frequency and orientation of the stimuli. The optimal PB model assumes that the participant used the rule depicted in Figure 1 as the solid line. This model has one free parameter representing the rule application variability (i.e., perceptual and criterial noise or σ^2). The second procedural-based model was the general linear classifier (GLC), which also assumes that the participant's decision on each trial is based on a linear integration of information from both the spatial frequency and orientation dimensions, although the weighting given to the two dimensions may be unequal. This model has three parameters, including the slope and intercept of the linear bound, and the rule application variability (i.e., σ^2).

The second class of models was the *hypothesis testing* (HT) models, which assumed that the participant used a verbalizable rule when performing the task. Two conjunctive rule models were applied to the data, both of which assume that the participant based her categorization decision on both dimensions. Both models as-




Figure 2. Category structure for the experiment.

sumed that the participant set a criterion along the spatial frequency dimension that distinguished low from high frequency stimuli, and set a criterion along the orientation dimension that distinguished shallow from steep angled stimuli. Thus the model assumed that there were four response regions: low frequency/shallow angle, low frequency/steep angle, high frequency/shallow angle and high frequency/steep angle. The criterion along the spatial frequency dimension was a free parameter and the criterion along the orientation dimension was a free parameter. One model assumed that low frequency/steep angle stimuli were in category A and all other stimuli were in category B. The other model assumed that high frequency/shallow angle stimuli were in category B and all other stimuli were in category A. Both models also included a rule application variability parameter (i.e., σ^2), for a total of three parameters in each model. Two additional hypothesis-testing models that were applied were unidimensional models, one that assumes the participant categorized the stimuli based only on the spatial frequency dimension and another that assumes the participant categorized the stimuli based only on the orientation dimension. These two models had two free parameters, a decision criterion on one of the dimensions and σ^2 . A final model that was applied assumes that the participant responded randomly (RR). Akaike's (Akaike, 1974) information criterion (AIC) was used to determine the model that provided the best account of the data.

Statistical Analyses

Demographic and behavioral results across groups were analyzed using SPSS 18.0 statistical software package using two-tailed independent *t* tests. Task performance was examined by contrasting participants' accuracy (percent correct), response time (reaction time (RT)), and RT variability (mean standard deviation for each block) across the entire 320 trials in 80 trial blocks using a 2 (group: AN vs. CW) \times 4 (blocks 1–4) mixed-design ANOVAs.

Results

Clinical Scales

The mean subscale and total scores for AN and CW on the BDI, TCI, EDI-2, and the SPSRQ are provided in Table 3. AN patients scored higher on the BDI than CW participants ($p < .001$). For the TCI, AN patients scored higher than the CW on the Harm Avoidance scales ($p < .05$). For the EDI-2, the AN group had significantly greater scores than the CW group on all subscales (all $ps < .01$). For the SPSRQ, the AN group had greater scores on both the SP and SR subscales (both $ps < .05$). There were no significant differences in scores on any clinical measures between restricting-type and binge/purging-type anorexia.

Task Accuracy Results

Accuracy results for the category learning task are shown in Figure 3. An ANOVA identified a main effect of group, $F(1, 38) = 4.5, p < .05, \eta_p^2 = .11$, with AN participants performing worse than the CW participants, and a main effect of block, $F(3, 114) = 10.6, p < .001, \eta_p^2 = .27$, with both AN and CW participants' performances improving across the trials. There was a trend for a

Table 3
Behavioral Data as Measured Through the Beck Depression Inventory (BDI), Temperament and Character Inventory (TCI), Eating Disorders Inventory 2 (EDI-2) Raw scores, and the Sensitivity to Punishment and Sensitivity to Reward (SPSRQ)

	CW		AN	
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>
BDI	1.42	1.7	26.7	11.4
TCI				
Novelty Seeking	20.1	5.2	16.5	6.2
Harm Avoidance	8.5	3.5	20.8	6.9
EDI-2				
Drive for Thinness	0.9	2.1	13.5	6.1
Bulimia	0.2	0.5	5.3	5.7
Body Dissatisfaction	2.2	2.7	14.0	9.4
Perfectionism	5.3	4.3	10.2	5.0
SPSRQ				
Sensitivity to Punishment	8.2	1.0	11.3	1.0
Sensitivity to Reward	5.4	0.7	13.1	1.1

Group X block interaction, $F(3, 114) = 2.4, p = .07, \eta_p^2 = .08$. Despite the lack of a significant Group X block interaction, follow-up *t* tests demonstrated that the two groups did not differ in either blocks 1 or 2 (both $ps > .17$), but the two groups did differ in blocks 3 and 4 (both $ps < .05; \eta_p^2 = .14$ for block 3, and $\eta_p^2 = .16$ for block 4). There was no difference in task performance in the subgroups of AN patients.

Task RT Results

In addition to accuracy, we also examined RT results in the CW and AN groups to help determine if other factors (e.g., attention and/or fatigue) might account for the accuracy differences described above. It was anticipated that if these other factors could account for the accuracy differences, then the AN group would likely have greater and more variable RTs than CW, particularly in the latter two blocks of trials where the accuracy levels differed. For mean RTs, the ANOVA did not identify a main effect of group, $F(1, 38) = 1.1, p > .05$, but there was a main effect of block, $F(1, 38) = 11.5, p < .01, \eta_p^2 = .24$, with both AN and CW participants' RTs becoming faster across the blocks. Importantly, there was no Group X block interaction, $F(3, 114) = 0.8, p > .05$, indicating that RTs did not differ between the groups across blocks. As for the RT standard deviations, there were no main effects of group, $F(1, 38) = 1.9, p > .05$, or block, $F(3, 114) = 0.58, p > .05$, and no Group X block interaction, $F(3, 114) = 0.63, p > .05$.

Task Performance Correlations With Clinical Data

To examine the relationship between performances on the category learning task and the clinical measures, a learning slope was computed by taking the final block accuracy and subtracting the first block accuracy for each participant. This learning slope was then correlated with the BDI and various subscales from the TCI and EDI-2 using Pearson correlations. The correlation coefficients for the TCI and EDI-2 are displayed in Table 4. For the TCI, the learning slope in the AN patients correlated significantly with the




Figure 3. Implicit learning task accuracy results (* $p < .05$).

Novelty Seeking subscale, such that patients with lower learning slopes (less learning) scored higher on this scale. For the SPSRQ, there was a significant positive correlation between learning slope and the SP subscale ($p < .05$) indicating that those AN patients with lower learning slopes were less sensitive to punishment. Learning slopes were not associated with the BDI, the included subscales of the EDI-2, lowest BMI, duration of illness, or age of disease onset. There were no significant correlations for the CW group between learning slope and any of the clinical measures.

Model Results

To determine whether groups differed in how they solved the categorization task in the last block of trials, we examined the number of participants in each group whose data were best fit by

Table 4
Correlation Coefficients for Learning Slope and the Clinical Measures * $p < .05$.

	CW	AN
BDI	-.14	.07
TCI		
Novelty Seeking	-.34	-.47*
Harm Avoidance	.32	.08
EDI-2		
Drive for Thinness	-.16	.08
Bulimia	.12	.11
Body Dissatisfaction	-.42	-.07
Perfectionism	.37	.11
SPSRQ		
Sensitivity to Punishment	.04	.44*
Sensitivity to Reward	-.06	.20

one of the PB model (i.e., the optimal model or the general linear classifier model) versus the number of participants whose data were best fit by one of the HT models described above (i.e., the conjunctive or unidimensional models). If one of the PB models best accounted for a participant's data, it would indicate that they based their decision on an implicit integration of the two stimulus dimensions, whereas if one of the HT model best accounted for a participant's data, it would indicate that this individual based their decision on a verbalizable rule. Finally, if the RR model best accounted for a participant's data it would indicate that they were responding randomly.

The results of these analyses for the final block indicated that, for the CW group, nine data sets were best fit by a PB model, nine were best fit by an HT model, and one was best fit by the RR model. For the AN group, nine data sets were best fit by a PB model, seven were best fit by an HT model, and five was best fit by the RR model. These frequencies did not differ based on a χ^2 test ($p = .22$). Thus, although there were differences in accuracy rates between the CW and AN groups, the two groups did not differ in terms of the frequency of participants who used an implicit (PB), verbalizable (HT), or random responding approach.

To further determine whether the model-based subgroups differed, we contrasted accuracy rates in the final block for the CW and AN participants who used either a PB or HT approach. Final block accuracy rates were used because these were the data on which the model analyses were conducted. These accuracy rates are shown in Figure 4. *T* tests demonstrated that the CW participants whose data were best fit by a PB model performed more accurately than those AN participants whose data were best fit by a PB model ($p < .05$). In contrast, accuracy in the final block for the CW and AN participants whose data were best fit by an HT model did not differ significantly ($p = .75$). Thus, accuracy rates differed between CW and AN participants who used an implicit approach to learning the task, whereas the groups did not differ in those participants who used a verbalizable approach. Within group comparisons between participants who used a PB or HT approach indicated that the AN PB users tended to perform worse than AN HT users (see Figure 4), although this difference was only a trend




Figure 4. Accuracy results for the Hypothesis Tested (HT) versus Procedural Based (PB) method of learning (* $p < .05$).

($p = .10$), whereas there was no difference between PB and HT users in the CW group ($p = .36$).

Finally, to determine if the AN participants whose final block data were best fit by a PB or HT model differed clinically, we contrasted these subgroups on the clinical variables and measures. The only difference that emerged was that the AN participants whose data was best fit by a PB model had larger scores on the Perfectionism subscale of the EDI-2 as compared to those participants whose data was best fit by an HT model (13.3 vs. 7.6; $p < .01$).

Discussion

To our knowledge, this is the first study to examine implicit category learning in patients with AN. The study has several key findings. First, individuals with AN performed worse on the implicit learning task than normal weight, age-matched controls, thus providing evidence for a deficit in implicit learning in ill AN. Second, as indicated by our model-based analyses, AN participants who used the task appropriate strategy still performed significantly worse than CW participants, indicating that AN women who learned the appropriate task approach were still impaired in learning, and poor strategy selection, *per se*, could not account for the learning deficit. Third, certain temperament and personality characteristics were associated with impaired category learning, such as novelty seeking and reinforcement biases (sensitivity to punishment).

Previous work has strongly indicated a role of dopamine in reward-based learning (Schultz, 1998, 2002) and recent research suggests that AN may be associated with alterations in the dopamine system (Fladung et al., 2010; Frieling et al., 2010; Kaye et al., 1984). The finding that AN patients are impaired in implicit category learning, which is believed to rely heavily on reward-based learning, provides support for a possible deficiency in dopamine functioning in AN patients, and is also in line with other studies that have suggested that certain forms of learning in AN are associated with deficits in the reward-based learning system (Lawrence et al., 2003). This possibility is also consistent with previous studies that other patient groups who have involvement of the dopamine system (such as patients with Parkinson's disease) are also impaired in implicit category learning (Buytenhuijs et al., 1994; Euteneuer et al., 2009; Filoteo, Maddox, Salmon, & Song, 2007; Price, 2005).

Although it is reasonable to speculate that dopamine abnormalities account for AN patients' deficits on the implicit category learning task, other possibilities must be considered. Previous studies have consistently identified deficits in AN patients in two other cognitive domains, cognitive set shifting and visual central coherence (Lopez, Tchanturia, Stahl, & Treasure, 2008; Lopez, Tchanturia, Stahl, Booth, Holliday, & Treasure, 2007; Roberts, Tchanturia, & Treasure, 2010; Southgate, Tchanturia, & Treasure, 2008; Tchanturia et al., 2004; Tchanturia et al., 2004; Tchanturia, Morris, Surguladze, & Treasure, 2002; Tenconi et al., 2010; Zastrow et al., 2009). Cognitive set shifting refers to those processes needed to switch from one cognitive set to another during the course of a task. Visual central coherence, on the other hand, refers to the ability to integrate and process an entire visual scene, as opposed to focusing on the component processes of the visual scene. Deficits in either of these two cognitive processes might also

account for AN patients' deficits in implicit category learning. Specifically, in the initial stages of performing an implicit category learning task, participants are thought to first attempt verbalizable rules in which the rule is often based on a single dimension, and only later do participants shift to a more accurate implicit rule (Ashby et al., 1998). In this study, if AN patients are impaired in cognitive set shifting, they might tend to continue to attempt to implement a verbal rule rather than switching and adopting a more accurate implicit rule, which would lead to a learning deficit. However, a similar proportion of study participants in each group adopted the implicit approach, suggesting that the AN group did not differentially stay with a rule based approach. The second possible explanation of the present findings may be that AN patients failed to see the gestalt of the stimuli because of a deficit in visual central coherence, and as such, they were unable to correctly learn to categorize the stimuli because they could not integrate the two stimulus dimensions. This might be consistent with previous findings that AN individuals had altered strategies and abilities to copy complex figures. Such a deficit might result in the AN patients focusing on only one dimension of the stimulus despite the fact that optimal responding required learning to categorize the stimuli based on both dimensions. Although both are plausible alternative explanations, it should be pointed out that even AN patients who adopted an implicit rule (as demonstrated by the model-based analyses) were impaired in learning the task relative to the controls. Those AN patients who used such an approach were able to shift cognitive set and integrate the stimulus dimensions, but nevertheless demonstrated impaired learning. Thus, although deficits in either cognitive set shifting or visual central coherence might account for the AN patients' deficits, it appears that a deficit in reward-based learning (possibly secondary to dopamine deficiencies) might best account for our findings.

Other findings from the current study supporting the notion that a deficit in reward-based learning accounts for AN patients' deficits was the association between poor performance in implicit category learning and personality assessments. In particular, impaired implicit category learning was associated with greater scores on the Novelty Seeking TCI subscale. Greater scores on Novelty Seeking are thought to reflect an individual's propensity to be more exploratory, impulsive and disorderly. Another finding was that poor category learning in AN patients was associated with less sensitivity to punishment based on the SPSRQ. Taken together, these results indicate that those individuals with AN who dislike rules and are less sensitive to punishment will be more likely to be impaired in implicit category learning.

The notion that AN patients are impaired in reward-based processes is not entirely new and has been the focus of several recent neurobiological and behavioral reviews (Harrison, O'Brien, Lopez, & Treasure, 2010; Kaye, Fudge, & Paulus, 2009; Keating, 2010; Scheurink, Boersma, Nergardh, & Sodersten, 2010). Claes and colleagues examined sensitivity to punishment as measured by the Behavioral Inhibition System subscale (Carver & White, 1994) and found that restricting subtype AN were more sensitive to punishment than purging subtype AN and healthy controls (Claes, Nederkoorn, Vandereycken, Guerrieri, & Vertommen, 2006). A recent study from our group using the SPSRQ found that restricting as well as binge eating/purging type AN had increased sensitivity to punishment and reward compared to age-matched healthy controls (Jappe et al., 2011). The present study also found similar

increased sensitivity to punishment and reward in AN as measured by the SPSRQ, but interestingly category learning performance was only associated with alterations in punishment sensitivity. AN individuals have traditionally been associated with perfectionism and high motivation to perform well (Bachner-Melman et al., 2007; Castro-Fornieles et al., 2007; Kaye, 2008; Maia et al., 2009; Nilsson, Sundbom, Dundbom, & Hagglof, 2008; Peck & Lightsey, 2008; Wade et al., 2008) and it could be possible that fear of punishment or failure have the benefit of higher effort and better performance. Still, biologic abnormalities in AN might prevent that group from performing similar to controls in implicit learning and other cognitive tasks. Interestingly, only the AN with highest perfectionism scores used the correct, implicit strategy. This could indicate that only those subjects who put forth the most effort actually ended up using the procedural based learning approach, while this did not seem to be a necessary factor for the controls. Further study is needed to examine this finding given that it indicates that there are some personality traits that might in fact positively influence learning in AN.

While the nature of this relationship should be explored further, it is important to point out that this present study is one of the first to link alterations in feedback sensitivity to impaired implicit learning processes in AN.

There are several limitations to the present study that should be addressed. First the sample sizes and cultural homogeneity of our participants limit the power of our statistical analyses and the generalizability of our findings to the general AN population. Second, we provide no direct link between the proposed alteration in the dopamine system in AN patients and their observed impairment on the implicit category learning task. Future research will have to address this hypothesized relationship. Third, at present the clinical implications of our findings is unknown, but we hope that our results provide the impetus for exploring such cognitive and temperament associations and how they impact clinically relevant aspects of the disease (e.g., prevention and treatment). Fourth, we did not have a direct measure of attention and/or fatigue, factors that could have impacted performance and account for the accuracy differences. However, the AN and CW groups did not differ in mean or variability in RT, suggesting that attention or fatigue may not have accounted for the AN learning deficit. Finally, the present study did not characterize the observed category learning deficit relative to other cognitive factors that might have influenced performance, such as general intelligence, so the specificity of this deficit is not known. It will be important for future studies to examine learning deficits in the context of other, potential cognitive impairment in AN individuals.

References

- Akaike, H. (1974). A new look at the statistical model identification. *IEEE Transactions on Automatic Control*, *19*, 716–723. doi:10.1109/TAC.1974.1100705
- Anderluh, M. B., Tchanturia, K., Rabe-Hesketh, S., & Treasure, J. (2003). Childhood obsessive-compulsive personality traits in adult women with eating disorders: Defining a broader eating disorder phenotype. *American Journal of Psychiatry*, *160*, 242–247. doi:10.1176/appi.ajp.160.2.242
- APA. (Ed.). (2000). *Diagnostic & Statistical Manual of Mental Disorders: DSM-IV-TR*. Washington, DC: American Psychiatric Association.
- Ashby, F. G. (Ed.). (1992). *Multidimensional models of perception and cognition*. Hillsdale, NJ: Lawrence Erlbaum Associates, Inc.
- Ashby, F. G., Alfonso-Reese, L. A., Turken, A. U., & Waldron, E. M. (1998). A neuropsychological theory of multiple systems in category learning. *Psychological Review*, *105*, 442–481. doi:10.1037/0033-295X.105.3.442
- Ashby, F. G., & Gott, R. E. (1988). Decision rules in the perception and categorization of multidimensional stimuli. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *14*, 33–53. doi:10.1037/0278-7393.14.1.33
- Ashby, F. G., & Maddox, W. T. (1992). Complex decision rules in categorization - contrasting novice and experienced performance. *Journal of Experimental Psychology: Human Perception and Performance*, *18*, 50–71. doi:10.1037/0096-1523.18.1.50
- Bachner-Melman, R., Lerer, E., Zohar, A. H., Kremer, I., Elizur, Y., Nemanov, L., . . . Ebstein, R. P. (2007). Anorexia nervosa, perfectionism, and dopamine D4 receptor (DRD4). *American Journal of Medical Genetics Part B-Neuropsychiatric Genetics*, *144B*, 748–756. doi:10.1002/ajmg.b.30505
- Beck, A. T., Ward, C. H., Mendelson, M., Mock, J., & Erbaugh, J. (1961). An inventory for measuring depression. *Archives of General Psychiatry*, *4*, 561–571. doi:10.1001/archpsyc.1961.01710120031004
- Bosanac, P., Kurlender, S., Stojanovska, L., Hallam, K., Norman, T., McGrath, C., . . . Olver, J. (2007). Neuropsychological study of underweight and “weight-recovered” anorexia nervosa compared with bulimia nervosa and normal controls. *International Journal of Eating Disorders*, *40*, 613–621. doi:10.1002/eat.20412
- Brainard, D. H. (1997). Psychophysics software for use with MATLAB. *Spatial Vision*, *10*, 433–436.
- Bulik, C. M., Tozzi, F., Anderson, C., Mazzeo, S. E., Aggen, S., & Sullivan, P. F. (2003). The relation between eating disorders and components of perfectionism. *American Journal of Psychiatry*, *160*, 366–368. doi:10.1176/appi.ajp.160.2.366
- Buytenhuis, E. L., Berger, H. J., Van Spaendonck, K. P., Horstink, M. W., Borm, G. F., & Cools, A. R. (1994). Memory and learning strategies in patients with Parkinson’s disease. *Neuropsychologia*, *32*, 335–342. doi:10.1016/0028-3932(94)90135-X
- Carver, C. S., & White, T. L. (1994). Behavioral inhibition, behavioral activation, and affective responses to impending reward and punishment: The BIS/BAS Scales. *Journal of Personality and Social Psychology*, *67*, 319–333. doi:10.1037/0022-3514.67.2.319
- Castro-Fornieles, J., Gual, P., Lahortiga, F., Gila, A., Casula, V., Fuhrmann, C., . . . Toro, J. (2007). Self-oriented perfectionism in eating disorders. *International Journal of Eating Disorders*, *40*, 562–568. doi:10.1002/eat.20393
- Cavanagh, J. F., Frank, M. J., & Allen, J. J. (2011). Social stress reactivity alters reward and punishment learning. *Social Cognitive and Affective Neuroscience*, *6*, 311–320. doi:10.1093/scan/nsq041
- Chui, H. T., Christensen, B. K., Zipursky, R. B., Richards, B. A., Hanratty, M. K., Kabani, N. J., . . . Katzman, D. K. (2008). Cognitive function and brain structure in females with a history of adolescent-onset anorexia nervosa. *Pediatrics*, *122*, e426–37. doi:10.1542/peds.2008-0170
- Claes, L., Nederkoorn, C., Vandereycken, W., Guerrieri, R., & Vertommen, H. (2006). Impulsiveness and lack of inhibitory control in eating disorders. *Eating Behaviors*, *7*, 196–203. doi:10.1016/j.eatbeh.2006.05.001
- Cloninger, C. R., Svrakic, D. M., & Przybeck, T. R. (1993). A psychobiological model of temperament and character. *Archives of General Psychiatry*, *50*, 975–990. doi:10.1001/archpsyc.1993.01820240059008
- Euteneuer, F., Schaefer, F., Stuermer, R., Boucsein, W., Timmermann, L., Barbe, M. T., . . . Kalbe, E. (2009). Dissociation of decision-making under ambiguity and decision-making under risk in patients with Parkinson’s disease: A neuropsychological and psychophysiological study.

- Neuropsychologia*, 47, 2882–2890. doi:10.1016/j.neuropsychologia.2009.06.014
- Fassino, S., Abbate-Daga, G., Amianto, F., Leombruni, P., Boggio, S., & Rovera, G. G. (2002). Temperament and character profile of eating disorders: A controlled study with the Temperament and Character Inventory. *International Journal of Eating Disorders*, 32, 412–425. doi:10.1002/eat.10099
- Filoteo, J. V., & Maddox, W. T. (1999). Quantitative modeling of visual attention processes in patients with Parkinson's disease: Effects of stimulus integrality on selective attention and dimensional integration. *Neuropsychology*, 13, 206–222. doi:10.1037/0894-4105.13.2.206
- Filoteo, J. V., Maddox, W. T., & Davis, J. D. (2001). A possible role of the striatum in linear and nonlinear category learning: Evidence from patients with Huntington's disease. *Behavioral Neuroscience*, 115, 786–798. doi:10.1037/0735-7044.115.4.786
- Filoteo, J. V., Maddox, W. T., Salmon, D. P., & Song, D. D. (2005). Information-integration category learning in patients with striatal dysfunction. *Neuropsychology*, 19, 212–222. doi:10.1037/0894-4105.19.2.212
- Filoteo, J. V., Maddox, W. T., Salmon, D. P., & Song, D. D. (2007). Implicit category learning performance predicts rate of cognitive decline in nondemented patients with Parkinson's disease. *Neuropsychology*, 21, 183–192. doi:10.1037/0894-4105.21.2.183
- Fladung, A. K., Gron, G., Grammer, K., Herrnberger, B., Schilly, E., Grasteit, S., . . . von Wietersheim, J. (2010). A neural signature of anorexia nervosa in the ventral striatal reward system. *American Journal of Psychiatry*, 167, 206–212. doi:10.1176/appi.ajp.2009.09010071
- Frieling, H., Beyer, S., Kalb, R., Kornhuber, J., Demling, J., de Zwaan, M., & Bleich, S. (2010). Anorexia nervosa in a 56-year-old woman with a diagnosis of hyperlipidemia: A case report. *Australian and New Zealand Journal of Psychiatry*, 44, 492–493. doi:10.3109/00048671003614155
- Garner, D. M. (1991). *Eating Disorder Inventory-2: Professional Manual*. Odessa, FL: Psychological Assessment Resources.
- Gray, J. A. (1970). The psychophysiological basis of introversion-extraversion. *Behaviour Research and Therapy*, 8, 249–266. doi:10.1016/0005-7967(70)90069-0
- Grunwald, M., Ettrich, C., Krause, W., Assmann, B., Dahne, A., Weiss, T., & Gertz, H. J. (2001). Haptic perception in anorexia nervosa before and after weight gain. *Journal of Clinical and Experimental Neuropsychology*, 23, 520–529. doi:10.1076/jcen.23.4.520.1229
- Harrison, A., O'Brien, N., Lopez, C., & Treasure, J. (2010). Sensitivity to reward and punishment in eating disorders. *Psychiatry Research*, 177, 1–11. doi:10.1016/j.psychres.2009.06.010
- Hermans, D., Pieters, G., & Eelen, P. (1998). Implicit and explicit memory for shape, body weight, and food-related words in patients with anorexia nervosa and nondieting controls. *Journal of Abnormal Psychology*, 107, 193–202. doi:10.1037/0021-843X.107.2.193
- Holliday, J., Tchanturia, K., Landau, S., Collier, D., & Treasure, J. (2005). Is impaired set-shifting an endophenotype of anorexia nervosa? *American Journal of Psychiatry*, 162, 2269–2275. doi:10.1176/appi.ajp.162.12.2269
- Jappe, L. M., Frank, G. K. W., Shott, M. E., Rollin, M. D. H., Pryor, T., Hagman, J. O., . . . Davis, E. (2011). Heightened sensitivity to reward and punishment in anorexia nervosa. *International Journal of Eating Disorders*, 44, 317–324. doi:10.1002/eat.20815
- Johansson, L., Ghaderi, A., & Andersson, G. (2005). Stroop interference for food- and body-related words: A meta-analysis. *Eating Behaviors*, 6, 271–281. doi:10.1016/j.eatbeh.2004.11.001
- Johansson, L., Ghaderi, A., Hallgren, M., & Andersson, G. (2008). Implicit memory bias for eating- and body appearance-related sentences in eating disorders: An application of Jacoby's white noise task. *Cognitive Behaviour Therapy*, 37, 135–145. doi:10.1080/16506070701664821
- Jones, B. P., Duncan, C. C., Brouwers, P., & Mirsky, A. F. (1991). Cognition in eating disorders. *Journal of Clinical and Experimental Neuropsychology*, 13, 711–728. doi:10.1080/01688639108401085
- Kaye, W. (2008). Neurobiology of anorexia and bulimia nervosa. *Physiology and Behavior*, 94, 121–135. doi:10.1016/j.physbeh.2007.11.037
- Kaye, W. H., Ebert, M. H., Gwirtsman, H. E., & Weiss, S. R. (1984). Differences in brain serotonergic metabolism between nonbulimic and bulimic patients with anorexia nervosa. *American Journal of Psychiatry*, 141, 1598–1601.
- Kaye, W. H., Fudge, J. L., & Paulus, M. (2009). New insights into symptoms and neurocircuit function of anorexia nervosa. *Nature Reviews Neuroscience*, 10, 573–584. doi:10.1038/nrn2682
- Keating, C. (2010). Theoretical perspective on anorexia nervosa: The conflict of reward. *Neuroscience and Behavioral Reviews*, 34, 73–79.
- Kingston, K., Szmukler, G., Andrewes, D., Tress, B., & Desmond, P. (1996). Neuropsychological and structural brain changes in anorexia nervosa before and after refeeding. *Psychological Medicine*, 26, 15–28. doi:10.1017/S0033291700033687
- Lauer, C. J., Gorzewski, B., Gerlinghoff, M., Backmund, H., & Zihl, J. (1999). Neuropsychological assessments before and after treatment in patients with anorexia nervosa and bulimia nervosa. *Journal of Psychiatry Research*, 33, 129–138. doi:10.1016/S0022-3956(98)00020-X
- Lawrence, A. D., Dowson, J., Foxall, G. L., Summerfield, R., Robbins, T. W., & Sahakian, B. J. (2003). Impaired visual discrimination learning in anorexia nervosa. *Appetite*, 40, 85–89. doi:10.1016/S0195-6663(02)00138-1
- Lena, S. M., Fiocco, A. J., & Leyenaar, J. K. (2004). The role of cognitive deficits in the development of eating disorders. *Neuropsychology Review*, 14, 99–113. doi:10.1023/B:NERV.0000028081.40907.de
- Lopez, C., Stahl, D., & Tchanturia, K. (2010). Estimated intelligence quotient in anorexia nervosa: A systematic review and meta-analysis of the literature. *Annals of General Psychiatry*, 9, 40. doi:10.1186/1744-859X-9-40
- Lopez, C., Tchanturia, K., Stahl, D., & Treasure, J. (2008). Central coherence in eating disorders: A systematic review. *Psychological Medicine*, 38, 1393–1404. doi:10.1017/S0033291708003486
- Lopez, C., Tchanturia, K., Stahl, D., Booth, R., Holliday, J., & Treasure, J. (2007). An examination of the concept of central coherence in women with anorexia nervosa. *International Journal of Eating Disorders*, 41, 143–152. doi:10.1002/eat.20478
- Maddox, W. T. (1999). On the dangers of averaging across observers when comparing decision bound models and generalized context models of categorization. *Perception and Psychophysics*, 61, 354–374. doi:10.3758/BF03206893
- Maddox, W. T. (2001). Separating perceptual processes from decisional processes in identification and categorization. *Perception and Psychophysics*, 63, 1183–1200. doi:10.3758/BF03194533
- Maddox, W. T., & Ashby, F. G. (1993). Comparing decision bound and exemplar models of categorization. *Perception and Psychophysics*, 53, 49–70. doi:10.3758/BF03211715
- Maddox, W. T., & Ashby, F. G. (1996). Perceptual separability, decisional separability, and the identification-speeded classification relationship. *Journal of Experimental Psychology: Human Perception and Performance*, 22, 795–817. doi:10.1037/0096-1523.22.4.795
- Maddox, W. T., & Ashby, F. G. (2004). Dissociating explicit and procedural-learning based systems of perceptual category learning. *Behavioural Processes*, 66, 309–332. doi:10.1016/j.beproc.2004.03.011
- Maddox, W. T., Ashby, F. G., & Bohil, C. J. (2003). Delayed feedback effects on rule-based and information-integration category learning. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 29, 650–662. doi:10.1037/0278-7393.29.4.650
- Maddox, W. T., & Filoteo, J. V. (2001). Striatal contributions to category learning: Quantitative modeling of simple linear and complex nonlinear rule learning in patients with Parkinson's disease.

- Journal of the International Neuropsychological Society*, 7, 710–727. doi:10.1017/S1355617701766076
- Maddox, W. T., Filoteo, J. V., Delis, D. C., & Salmon, D. P. (1996). Visual selective attention deficits in patients with Parkinson's disease: A quantitative model-based approach. *Neuropsychology*, 10, 197–218. doi:10.1037/0894-4105.10.2.197
- Maddox, W. T., Filoteo, J. V., & Huntington, J. R. (1998). Effects of stimulus integrality on visual attention in older and younger adults: A quantitative model-based analysis. *Psychology and Aging*, 13, 472–485. doi:10.1037/0882-7974.13.3.472
- Maddox, W. T., Filoteo, J. V., Lauritzen, J. S., Connally, E., & Hejl, K. D. (2005). Discontinuous categories affect information-integration but not rule-based category learning. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 31, 654–659. doi:10.1037/0278-7393.31.4.654
- Maia, B. R., Soares, M. J., Gomes, A., Marques, M., Pereira, A. T., Cabral, A., . . . Macedo, A. (2009). Perfectionism in obsessive-compulsive and eating disorders. *Revista Brasileira De Psiquiatria*, 31, 322–327. doi:10.1590/S1516-44462009005000004
- Mathias, J. L., & Kent, P. S. (1998). Neuropsychological consequences of extreme weight loss and dietary restriction in patients with anorexia nervosa. *Journal of Clinical and Experimental Neuropsychology*, 20, 548–564. doi:10.1076/jcen.20.4.548.1476
- Miyachi, S., Hikosaka, O., Miyashita, K., Karadi, Z., & Rand, M. K. (1997). Differential roles of monkey striatum in learning of sequential hand movement. *Experimental Brain Research*, 115, 1–5. doi:10.1007/PL00005669
- Nakamura, K., & Hikosaka, O. (2006). Role of dopamine in the primate caudate nucleus in reward modulation of saccades. *Journal of Neuroscience*, 26, 5360–5369. doi:10.1523/JNEUROSCI.4853-05.2006
- Nilsson, K., Sundbom, E., & Hagglof, B. (2008). A longitudinal study of perfectionism in adolescent onset anorexia nervosa-restricting type. *European Eating Disorders Review*, 16, 386–394. doi:10.1002/erv.850
- Nomura, E. M., Maddox, W. T., Filoteo, J. V., Ing, A. D., Gitelman, D. R., Parrish, T. B., . . . Reber, P. J. (2007). Neural correlates of rule-based and information-integration visual category learning. *Cerebral Cortex*, 17, 37–43. doi:10.1093/cercor/bhj122
- Ohrmann, P., Kersting, A., Suslow, T., Lalee-Mentzel, J., Donges, U. S., Fiebig, M., . . . Pfeleiderer, B. (2004). Proton magnetic resonance spectroscopy in anorexia nervosa: Correlations with cognition. *Neuroreport*, 15, 549–553. doi:10.1097/00001756-200403010-00033
- Palazidou, E., Robinson, P., & Lishman, W. A. (1990). Neuroradiological and neuropsychological assessment in anorexia nervosa. *Psychological Medicine*, 20, 521–527. doi:10.1017/S0033291700017037
- Peck, L. D., & Lightsey, O. R. (2008). The eating disorders continuum, self-esteem, and perfectionism. *Journal of Counseling and Development*, 86, 184–192.
- Price, A. L. (2005). Cortico-striatal contributions to category learning: Dissociating the verbal and implicit systems. *Behavioral Neuroscience*, 119, 1438–1447. doi:10.1037/0735-7044.119.6.1438
- Roberts, M. E., Tchanturia, K., Stahl, D., Southgate, L., & Treasure, J. (2007). A systematic review and meta-analysis of set-shifting ability in eating disorders. *Psychological Medicine*, 37, 1075–1084. doi:10.1017/S0033291707009877
- Roberts, M. E., Tchanturia, K., & Treasure, J. L. (2010). Exploring the neurocognitive signature of poor set-shifting in anorexia and bulimia nervosa. *Journal of Psychiatric Research*, 44, 964–970. doi:10.1016/j.jpsychires.2010.03.001
- Scheurink, A. J., Boersma, G. J., Nergardh, R., & Sodersten, P. (2010). Neurobiology of hyperactivity and reward: Agreeable restlessness in anorexia nervosa. *Physiology and Behavior*, 100, 490–495. doi:10.1016/j.physbeh.2010.03.016
- Schultz, W. (1998). Predictive reward signal of dopamine neurons. *Journal of Neurophysiology*, 80, 1–27.
- Schultz, W. (2002). Getting formal with dopamine and reward. *Neuron*, 36, 241–263. doi:10.1016/S0896-6273(02)00967-4
- Seed, J. A., Dixon, R. A., McCluskey, S. E., & Young, A. H. (2000). Basal activity of the hypothalamic-pituitary-adrenal axis and cognitive function in anorexia nervosa. *European Archives of Psychiatry and Clinical Neuroscience*, 250, 11–15. doi:10.1007/PL00007533
- Seed, J. A., McCue, P. M., Wesnes, K. A., Dahabra, S., & Young, A. H. (2002). Basal activity of the HPA axis and cognitive function in anorexia nervosa. *International Journal of Neuropsychopharmacology*, 5, 17–25. doi:10.1017/S146114570100270X
- Shafraan, R., Lee, M., Cooper, Z., Palmer, R. L., & Fairburn, C. G. (2007). Attentional bias in eating disorders. *International Journal of Eating Disorders*, 40, 369–380. doi:10.1002/eat.20375
- Shohamy, D., Myers, C. E., Onlaor, S., & Gluck, M. A. (2004). Role of the basal ganglia in category learning: How do patients with Parkinson's disease learn? *Behavioral Neuroscience*, 118, 676–686. doi:10.1037/0735-7044.118.4.676
- Southgate, L., Tchanturia, K., & Treasure, J. (2008). Information processing bias in anorexia nervosa. *Psychiatry Research*, 160, 221–227. doi:10.1016/j.psychres.2007.07.017
- Steinglass, J. E., Walsh, B. T., & Stern, Y. (2006). Set shifting deficit in anorexia nervosa. *Journal of the International Neuropsychological Society*, 12, 431–435. doi:10.1017/S1355617706060528
- Sullivan, P. F. (1995). Mortality in anorexia nervosa. *American Journal of Psychiatry*, 152, 1073–1074.
- Szmukler, G. I., Andrewes, D., Kingston, K., Chen, L., Stargatt, R., & Stanley, R. (1992). Neuropsychological impairment in anorexia nervosa: Before and after refeeding. *Journal of Clinical and Experimental Neuropsychology*, 14, 347–352. doi:10.1080/01688639208402834
- Tchanturia, K., Anderluh, M. B., Morris, R. G., Rabe-Hesketh, S., Collier, D. A., Sanchez, P., & Treasure, J. L. (2004). Cognitive flexibility in anorexia nervosa and bulimia nervosa. *Journal of the International Neuropsychological Society*, 10, 513–520. doi:10.1017/S1355617704104086
- Tchanturia, K., Morris, R. G., Anderluh, M. B., Collier, D. A., Nikolaou, V., & Treasure, J. (2004). Set shifting in anorexia nervosa: An examination before and after weight gain, in full recovery and relationship to childhood and adult OCPD traits. *Journal of Psychiatry Research*, 38, 545–552. doi:10.1016/j.jpsychires.2004.03.001
- Tchanturia, K., Morris, R. G., Surguladze, S., & Treasure, J. (2002). An examination of perceptual and cognitive set shifting tasks in acute anorexia nervosa and following recovery. *Eat and Weight Disorders*, 7, 312–315.
- Tchanturia, K., Whitney, J., & Treasure, J. (2006). Can cognitive exercises help treat anorexia nervosa? *Eat and Weight Disorders*, 11, e112–6.
- Tenconi, E., Santonastaso, P., Degortes, D., Bosello, R., Titton, F., Mapelli, D., & Favaro, A. (2010). Set-shifting abilities, central coherence, and handedness in anorexia nervosa patients, their unaffected siblings and healthy controls: Exploring putative endophenotypes. *The World Journal of Biological Psychiatry*, 11, 813–823.
- Torrubia, R., Avila, C., Molto, J., & Caseras, X. (2001). The Sensitivity to Punishment and Sensitivity to Reward Questionnaire (SPSRQ) as a measure of Gray's anxiety and impulsivity dimensions. *Personality and Individual Differences*, 31, 837–862. doi:10.1016/S0191-8869(00)00183-5
- Wächter, T., Lungu, O. V., Liu, T., Willingham, D. T., & Ashe, J. (2009). Differential effect of reward and punishment on procedural learning. *Journal of Neuroscience*, 29, 436–443. doi:10.1523/JNEUROSCI.4132-08.2009
- Wade, T. D., Tiggemann, M., Bulik, C. M., Fairburn, C. G., Wray, N. R., & Martin, N. G. (2008). Shared temperament risk factors for anorexia nervosa: A twin study. *Psychosomatic Medicine*, 70, 239–244. doi:10.1097/PSY.0b013e31815c40f1
- Wagner, A., Barbarich-Marsteller, N. C., Frank, G. K., Bailer, U. F.,

- Wonderlich, S. A., Crosby, R. D., . . . Kaye, W. H. (2006). Personality traits after recovery from eating disorders: Do subtypes differ? *International Journal of Eating Disorders, 39*, 276–284. doi: 10.1002/eat.20251
- Zastrow, A., Kaiser, S., Stippich, C., Walther, S., Herzog, W., Tchanturia, K., . . . Friederich, H. C. (2009). Neural correlates of impaired cognitive-behavioral flexibility in anorexia nervosa. *American Journal of Psychiatry, 166*, 608–616. doi:10.1176/appi.ajp.2008.08050775
- Zeithamova, D., & Maddox, W. T. (2006). Dual-task interference in perceptual category learning. *Memory & Cognition, 34*, 387–398. doi: 10.3758/BF03193416

Received April 19, 2011

Revision received October 5, 2011

Accepted November 21, 2011 ■